

## Features

### Weight

22 lbs (10 kg)

### Ingredients

calcium carbonate, monocalcium phosphate, salt, magnesium oxide, sodium sulfate, magnesium sulfate, wheat bran.

### Minimum guaranteed analysis

21,0 % Calcium (Ca)  
4,0 % Phosphore (P)  
7,5 % Salt  
3,5 % Sodium (Na)  
2,5 % Magnesium (Mg)

### Hardness level

Hard

### Recommended rate

1 PECKstone for every 500 birds

## Advantages

Aggressive pecking behavior: a common and serious welfare and economic problem. Newer housing systems may increase feather pecking, incidences of skeletal injuries, and mortality. 55-80% of producers report signs of feather pecking. Prevention is ideal and it's critical to intervene immediately to prevent further damage.

- A unique enrichment tool **designed specifically to prevent and reduce abnormal pecking behavior**
- Redirects aggressive pecking behavior
- Supports natural beak wear (with or without beak trimming) for optimal beak conditioning
- Durable yet destructible, an enticing pecking target
- **All natural and organic**
- Easy and practical for any cage-free operation
- Enhances animal welfare
- Can last between 4 and 8 weeks
- 24 month expiration date

## How to use and place the PECKStones

POULTRY TYPE	USE AND PLACEMENT
Cage-free layers and breeders	<ul style="list-style-type: none"> <li>• Use from initial bird placement, or add anytime</li> <li>• Remove the PECKStone from the plastic tub</li> <li>• Turn the tub upside down and place on litter or scratch area</li> <li>• Place the PECKStone on top of the tub</li> </ul>
Broilers and pullets	<ul style="list-style-type: none"> <li>• Use from initial bird placement</li> <li>• Remove the PECKStone from the plastic tub</li> <li>• Place the PECKStone on litter or scratch area</li> </ul>



**PREVENT  
AND  
REDUCE  
PECKING**

**ONE STONE  
IDEAL FOR  
500 BIRDS**

**FOR ANY  
POULTRY  
FACILITIES**



For a short explanatory video,  
click on the image above